



P B 5818 - Patentaan 2  
2280 HV Rijswijk  
T +31 70 340 3016  
TX 31651 epo  
FAX +31 70 340 3016

Europäisches  
Patentamt

Zweigstelle  
in Den Haag  
Recherchen-  
abteilung

European  
Patent Office

Branch at  
The Hague  
Search  
division

Office européen  
des brevets

Département a  
La Haye  
Division de la  
recherche

Ben Rudolf 17. am 13.12.02

Ciba Specialty Chemicals Holding Inc.  
Patentabteilung,  
Klybeckstrasse 141  
4057 Basel  
SUISSE

PATENTS		LE 5
12. Dez. 2002		
PATA	PATH	SES
/	/	REM

Datum/Date

12.12.02

Zeichen/Ref./Réf.

BS/1-22739/CGM5/19

Anmeldung Nr./Application No./Demande n°/Patent Nr./Patent No./Brevet n°.

02405763.0-2109-

Anmelder/Applicant/Demandeur/Patentinhaber/Proprietor/Titulaire

Ciba Specialty Chemicals Holding Inc.

## COMMUNICATION

The European Patent Office herewith transmits as an enclosure the European search report for the above-mentioned European patent application.

If applicable, copies of the documents cited in the European search report are attached.

☐ Additional set(s) of copies of the documents cited in the European search report is (are) enclosed as well.

The following specifications given by the applicant have been approved by the Search Division:

☐ abstract

☒ title

☒ The abstract was modified by the Search Division and the definitive text is attached to this communication.

The following figure will be published together with the abstract: NONE

## REFUND OF THE SEARCH FEE

If applicable under Article 10 Rules relating to fees, a separate communication from the Receiving Section on the refund of the search fee will be sent later.





## ABSTRACT / ZUSAMMENFASSUNG / ABREGE

02405763.0

Method for the preparation of a comb copolymer comprising

a) polymerising in a first step one or more epoxy group containing monomers to obtain a polyether, wherein at least one monomer is of formula (I)

wherein L is a linking group selected from the group consisting of  $C_1$ - $C_{18}$ alkylene, phenylene, phenylene- $C_1$ - $C_{18}$ alkylene,  $C_1$ - $C_{18}$ alkylene-phenylene,  $C_1$ - $C_{18}$ alkylene-phenylene-oxy and  $C_5$ - $C_{12}$ cycloalkylene;

$R_p$  and  $R_q$  are independently tertiary bound  $C_4$ - $C_{28}$ alkyl groups which are unsubstituted or substituted by one or more electron withdrawing groups or by phenyl; or

$R_p$  and  $R_q$  together form a 5 or 6 membered heterocyclic ring which is substituted at least by 4  $C_1$ - $C_4$ alkyl groups and which may be interrupted by a further nitrogen or oxygen atom; and in a second step

b) adding to the polymer obtained in step a) at least one ethylenically unsaturated monomer, heating the resulting mixture to a temperature where cleavage of the nitroxylether bond occurs and radical polymerization starts;

and polymerizing to the desired degree.